arXiv:1601.06255v1 [math.DG] 23 Jan 2016

PROJECTIVE CLASSIFICATION OF JETS OF SURFACES IN $\mathbb{P}^4$

JORGE LUIZ DEOLINDO SILVA AND YUTARO KABATA

Abstract. We are interested in the local extrinsic geometry of smooth surfaces in 4-space, and classify jets of Monge forms by projective transformations according to $\mathcal{A}^3$-types of their central projections.

1. Introduction

We are concerned with the geometry of the contact of smooth surfaces in the projective space $\mathbb{R}P^4$ with their tangent lines. The contact is measured by types of map germs of *central projections* of the surfaces. In the present paper we classify jets of generic surfaces by projective transformations which preserve the geometry of the contacts of surfaces with their tangent lines.

For surfaces in $\mathbb{R}P^3$, Platonova [20, 21] completed the classification of generic surfaces by projective transformations. The classification of generic two parameter families of surfaces is done in [8] (see also [1, 10, 11, 12]). The study of surfaces in $\mathbb{R}P^4$ was proposed in [2] with relation to the classification of singularities which appear in central projections of generic surfaces in $\mathbb{R}^4$ by D. Mond [15, 16, 17]. However there have been no results about the classification of surfaces so far. This paper gives an answer to the proposal in [2] with the complete classification list of jets of generic surfaces in $\mathbb{R}P^4$ by projective transformations.

On the other hand, the study of geometric aspects of surfaces in 4-space is a relatively new subject and has a lot of analogy to the study of surfaces in 3-space which have been investigated in [5, 6, 13, 14, 18].

Let M be a smooth surface embedded in $\mathbb{R}^4 \subset \mathbb{R}P^4$ containing the origin of $\mathbb{R}^4$ where $\mathbb{R}^4$ is identified with the open subset $\{[x; y; z; w; 1]\} \subset \mathbb{R}P^4$. We write $(z, w) = f(x, y) = (f_1(x, y), f_2(x, y))$ as the Monge form of M at the origin where $f_i(0, 0) = df_i(0, 0) = 0$ for $i = 1, 2$. Two jets of surfaces at some points are said to be *projective equivalent* if there is a projective transformation on $\mathbb{R}P^4$ sending one to the other. Our result is the following.

Theorem 1. *There is an open everywhere dense subset $\mathcal{O}$ of the space of compact smooth surfaces M in $\mathbb{R}P^4$ such that the germ at each point on M in $\mathcal{O}$ is projectively equivalent to a germ with the k-jet of the Monge form of one of the cases in Table 1.*

Observe that the last column of Table 1 means the types of central projections of corresponding surfaces from view points on asymptotic lines. In section 2, we briefly explain the stratification of the space of jets of Monge forms induced by the stratification of jets of germs of central projections, and review the stratification of the 3-jet space of Monge forms induced from the $\mathcal{A}^3$-stratification of central projection germs which was originally done in Ph. D thesis of Mond [15]. In Section 3 we obtain simple normal forms of jets of Monge forms that represent each

Type	Normal form	Condition	cod	Proj.
Π_E	$(x^2-y^2+y^2(\phi_1+\phi_2), xy+\psi_4)$		0	$-$
Π_S	$(x^2+y^3+y\phi_3, y^2+\alpha x^3+x\psi_3)$	$\alpha\neq 0$	0	S
Π_B	$(x^2+y^3+y\phi_3, y^2+x\psi_3)$	$-$	1	S,B
Π_{2B}	$(x^2+y\phi_3, y^2+x\psi_3)$	$-$	2	B
Π_H	$(x^2+\beta xy^2+y^3+y\phi_3, xy+x\psi_3)$	$-$	1	H
Π_P	$(x^2+xy^2+\lambda y^4, xy+\gamma y^3+\psi_4)$	$\gamma,\Lambda\neq 0$	2	P
Π_I^+	$(x^2+y^2+k_1x^2y+y\phi_3, \psi_3+\psi_4)$	$b_{30}-b_{12}\neq 0$	2	S,B
Π_I^-	$(xy+k_2x^3+\phi_4, \psi_3+\psi_4)$	$b_{30},b_{03}\neq 0, a_{22}=0$	2	S,B,H

TABLE 1. Strata of codimension ≤ 2 in the space of 4-jets of Monge forms corresponding to $\mathcal{A}^3$-types of central projections on asymptotic lines. Here $\phi_s=\sum_{i+j=s}a_{ij}x^iy^j, \psi_s=\sum_{i+j=s}b_{ij}x^iy^j$, $\alpha,\beta,\gamma,\lambda,k_1,k_2,a_{ij},b_{ij}\in\mathbb{R}$ are moduli parameters and $\Lambda=6\gamma^2+4\lambda-15\gamma+5$. Surface germs of Π_E-type do not have asymptotic lines.

stratum given in the previous section by projective transformations, and give the proof of Theorem 1.

Remark 1.1. *Our normal forms in Table 1 contain a lot of moduli parameters including coefficients of higher order terms of degree grater than* 4*. They must be interpreted as some projective differential invariants. For example, when we look at the $\mathcal{A}$-types of the central projection of the Π_S-type surfaces germs, it is observed that the central projection from a view point on the asymptotic line is $\mathcal{A}$-equivalent to $P_3(c):(x,xy^2+cy^4,xy+y^3)$ where c is a moduli parameter, and $\frac{\lambda}{\gamma}=c$. The first author* [7] *found also that γ and λ are expressed by combinations of some cross-ratio invariants and they determines the topological type of BDE (binary differential equations) of asymptotic curves.*

Acknowledgements: We would like to thank Takashi Nishimura and Farid Tari for organizing the JSPS-CAPES no.002/14 bilateral project in 2014-2016. The authors are supported by the project for their stays in ICMC-USP and Hokkaido University, respectively. The first author thanks also the FAPESP no.2012/ 00066-9 to support part of this work. We are also very grateful to Farid Tari and Toru Ohmoto for their supervisions.

2. The stratification of the 3-jet space of Monge forms

Mond, in Chapter III of his Ph.D thesis [15], stratified the jet space of Monge forms according to $\mathcal{A}$-types of germs of central projections. In this section we explain the way to stratify the jet space of Monge forms according to types of central projections, and review Mond's stratification for the 3-jet space of Monge forms.

Take a surface M in $\mathbb{R}^4\subset\mathbb{R}P^4$ with the Monge form $(z,w)=f(x,y)=(f_1(x,y),f_2(x,y))$. Let V_ℓ denote the space of polynomials in x,y of degree greater than 1 and less than or equal to ℓ. Our aim here is to obtain a stratification of the ℓ-jet space of Monge forms $V_\ell\times V_\ell$ which is induced from the $\mathcal{A}^\ell$-stratification of $J^\ell(2,3)$ obtained in [15, 16] as follows.

Type	Normal form	cod
hyperbolic	(x^2, y^2)	0
elliptic	$(x^2 - y^2, xy)$	0
parabolic	(x^2, xy)	1
inflection	$(x^2 + y^2, 0)$ or $(xy, 0)$	2
degenerate inflection	$(x^2, 0)$	3
degenerate inflection	$(0, 0)$	4

TABLE 2. The classification of $J^2(2,2)$ (which is equal to the 2-jet space of Monge forms $f = (f_1, f_2)$) by $GL(2)\times GL(2)$-actions given by Gibson in [9].

Consider a point $p \in \mathbb{R}P^4$, which is sometimes called *a view point*, not lying on M and define $\pi_p : \mathbb{R}P^4 - \{p\} \to \mathbb{R}P^3$ as the canonical projection which associates $x \in \mathbb{R}P^4 - \{p\}$ to the line generated by $x - p$. *The central projection of the surface M from p* is given by the composite map

$$\varphi_{p,M} := \pi_p \circ \iota : M \to \mathbb{R}P^3$$

(see also [8]).

We denote the central projection of the surface germ expressed in Monge form f from a view point p by $\varphi_{p,f}$. We stratify $V_\ell \times V_\ell$ by the difference of $\mathcal{A}^\ell$-types of $j^\ell\varphi_{p,f}$ for a view point $p \in \mathbb{R}P^4 - M$ and the ℓ-jet of a Monge form $j^\ell f \in V_\ell \times V_\ell$. $\mathcal{A}^\ell$ means the equivalence of jets of map germs, i.e., two jets $j^\ell g, j^\ell h \in J^\ell(2,3)$ are equivalent if and only if there exist jets of diffeomorphism germs σ, τ of the source and the target at the origins such that $j^\ell h = j^\ell(\tau\circ g\circ\sigma^{-1})$. Remark that even for the same ℓ-jet of the surface the central projection gives different $\mathcal{A}^\ell$-types depending on the view point. For example, $j^\ell\varphi_{p,f}$ is always regular type (i.e. equivalent to $(x, y, 0)$) if and only if p is outside the tangent plane of the surface germ; and gives a singularity if and only if p is on the tangent plane to the surface.

We say that a line on the tangent plane which goes through the origin in $\mathbb{R}^4$ is an *asymptotic line of the surface given in Monge form f* if $\varphi_{p,f}$ is equivalent to one of singularities worse than a crosscap (S_0-type) for all view points p on the line. In Section 3 we show that the classification of 2-jets of Monge forms by projective transformations coincides with the classification of $J^2(2,2)$ by $GL(2) \times GL(2)$-actions in Table 2, and each orbit is characterized by the number of asymptotic lines [5, 16]. For elliptic type with the form $j^2 f = (x^2 + y^2, xy)$, there are no asymptotic lines; for hyperbolic type with the form $j^2 f = (x^2, y^2)$, x and y-axis are two unique asymptotic lines; for parabolic type with the form $j^2 f = (x^2, xy)$, y-axis is a unique asymptotic line; for inflection type with the form $j^2 f = (x^2 + y^2, 0)$ or $(xy, 0)$, all lines on the tangent plane which go through the origin are asymptotic lines. Thus we study types of central projections from view points on the asymptotic lines, and divide strata in Table 2 into finer ones.

Define a smooth map, the Monge-Taylor map $\Theta : M \to V_\ell \times V_\ell$, which associates to each point in M the ℓ-jet of Monge form $f = (f_1, f_2)$ at the point. The following is a natural extension of Bruce's Theorem in [3].

Name	Normal form
S_0	(x, y^2, xy)
S	$(x, y^2, y^3 + x^2y)$ or (x, y^2, y^3)
B	(x, y^2, x^2y) or $(x, y^2, 0)$
H	(x, xy, y^3)
P	$(x, xy + y^3, xy^2)$
R	(x, xy, xy^2)
T	$(x, xy + y^3, 0)$
U	$(x, xy, 0)$

TABLE 3. $\mathcal{A}^3$-orbits of germs $\mathbb{R}^2, 0 \to \mathbb{R}^3, 0$ [15, 16].

Theorem 2. *Let $Z \subset V_\ell \times V_\ell$ be an $GL(2) \times GL(2)$-invariant submanifold. For generic surface M in $\mathbb{R}^4$, the Monge-Taylor map $\Theta : M \to V_\ell \times V_\ell$ is transverse to Z.*

Proof : The proof follows the same arguments in the proof of Theorem 1 in [3] (see also [6]). □

Since the strata are induced from $\mathcal{A}^\ell$-types of central projections, they are $GL(2) \times GL(2)$-invariant. By Theorem 2 we consider only strata with codimension at most 2. In this section, we study the stratification of 3-jets of Monge forms which is induced by the $\mathcal{A}^3$-orbits in [15, 16, 17] or their unions given in Table 3.

Write

$$f_1(x, y) = \sum_{i+j\geq 2} a_{ij}x^i y^i, \quad f_2(x, y) = \sum_{i+j\geq 2} b_{ij}x^i y^i.$$

Then $\varphi_{p,f}$ can be regarded as a map germ $\mathbb{R}^2, 0 \to \mathbb{R}^3$. Indeed, for $p = [a; b; c; d; 1] \in \mathbb{R}^4 - M$, we choose $a \neq 0$, then $\varphi_{p,f}$ is given by

$$\varphi_{p,f}(x, y) = \left(\frac{y - b}{x - a}, \frac{f_1(x, y) - c}{x - a}, \frac{f_2(x, y) - d}{x - a}\right).$$

On the other hand, if p is taken at infinity and written as $p = [a; b; c; d; 0]$, $\varphi_{p,f}$ is given by

$$\varphi_{p,f}(x, y) = (y - ux, f_1(x, y) - vx, f_2(x, y) - wx)$$

with $(u, v, w) = (\frac{b}{a}, \frac{c}{a}, \frac{d}{a})$ (see also [8]).

The following sums up the Propositions III. 2:2, 2:8, 2:14, 2:16 and 2:17 in [15].

Proposition 2.1. (Mond [15]) *For a surface germ in Monge form $(z, w) = f(x, y)$ we have the following.*

(i) *Suppose that $j^2 f = (x^2, y^2)$. Then:*

$$\begin{array}{rcl} j^3\varphi_{p,f} \sim S & \Leftrightarrow & a_{03} \neq 0 \ (\textit{resp. } b_{30} \neq 0) \\ j^3\varphi_{p,f} \sim B & \Leftrightarrow & a_{03} = 0 \ (\textit{resp. } b_{30} = 0) \end{array}$$

for p on the asymptotic line $x = 0$ (resp. $y = 0$).

(ii) *Suppose that* $j^2f = (x^2, xy)$. *Then:*

$$\begin{array}{lcl} j^3\varphi_{p,f} \sim H & \Leftrightarrow & a_{03} \neq 0 \\ j^3\varphi_{p,f} \sim P & \Leftrightarrow & a_{03} = 0 \text{ and } a_{12}, b_{03} \neq 0 \\ j^3\varphi_{p,f} \sim R & \Leftrightarrow & a_{03} = b_{03} = 0 \text{ and } a_{12} \neq 0 \\ j^3\varphi_{p,f} \sim T & \Leftrightarrow & a_{03} = a_{12} = 0 \text{ and } b_{03} \neq 0 \\ j^3\varphi_{p,f} \sim U & \Leftrightarrow & a_{03} = a_{12} = b_{03} = 0 \end{array}$$

for p *on the unique asymptotic line* $x = 0$.

(iii) *Suppose that* $j^2f = (x^2 + y^2, 0)$. *Then:*

$$j^3\varphi_{p,f} \sim S \text{ or } B$$

for any p *on the* xy*-plane.*

(iv) *Suppose that* $j^2f = (xy, 0)$. *Then:*

$$\begin{array}{lcl} j^3\varphi_{p,f} \sim S, B \text{ or } H & \Leftrightarrow & b_{30} \neq 0 \\ j^3\varphi_{p,f} \sim S, B \text{ or } P & \Leftrightarrow & b_{30} = 0 \text{ and } a_{30}, b_{21} \neq 0 \\ j^3\varphi_{p,f} \sim S, B \text{ or } R & \Leftrightarrow & a_{30} = b_{30} = 0 \text{ and } b_{21} \neq 0 \\ j^3\varphi_{p,f} \sim S, B \text{ or } T & \Leftrightarrow & a_{30} = b_{21} = 0 \text{ and } b_{30} \neq 0 \\ j^3\varphi_{p,f} \sim S, B \text{ or } U & \Leftrightarrow & a_{30} = b_{30} = b_{21} = 0 \end{array}$$

for any p *on the* xy*-plane.*

Proof : Statement (i). It is easy to check that the x and y-axes are asymptotic lines at the origin for M in Monge-form with $j^2f = (x^2, y^2)$. Suppose p is at the y-axis and written as $p = (0, a, 0, 0)$, then, by coordinate changes, we get

$$j^3\varphi_{p,f} \sim_{\mathcal{A}^3} (x, a(a_{21}a - 1)x^2y + a_{03}y^3, y^2).$$

If $a_{03} \neq 0$, $j^3\varphi_{p,f}(0)$ is of S-type, otherwise it is of B-type. If p is at infinity on the y-axis, we obtain

$$j^3\varphi_{p,f} \sim_{\mathcal{A}^3} (x, xy, a_{21}x^2y + a_{03}y^3).$$

Again, if $a_{03} \neq 0$, $j^3\varphi_{p,f}(0)$ is of S-type, otherwise it is of B-type. By exchanging x and y (also a_{ij} and b_{ij}), the case of p on the x-axis is follows similarly.

Statement (ii). Remark that y-axis is a unique asymptotic line at the origin when $j^2f(0) = (x^2, xy)$, hence we put $p = (0, a, 0, 0) \in \mathbb{R}^4 \subset \mathbb{R}P^4$ $(a \neq 0)$ and we get

$$j^3\varphi_{p,f} \sim_{\mathcal{A}^3} (x, a_{12}axy^2 + a_{03}y^3, xy + b_{03}y^3).$$

Statement (ii) for $p \in \mathbb{R}^4$ naturally follows, and the case p at infinity is similar.

Statement (iii). Let view point $p = (a, b, 0, 0) \in \mathbb{R}^4$ and $a \neq 0$, then we get

$$j^3\varphi_{p,f} \sim_{\mathcal{A}^3} (x, y^2, \xi_1x^2y + \xi_2y^3)$$

where ξ_1 and ξ_2 are homogeneous polynomials of degree 3 with variables a and b whose coefficients consist of a_{ij} and b_{ij}. Statement (iii) for $p = (a, b, 0, 0) \in \mathbb{R}^4$ where $a \neq 0$ naturally follows, and it is easily seen that the projection gives just S_1-type for view points $p = (0, b, 0, 0) \in \mathbb{R}^4$ where $b \neq 0$. The case p at infinity is similar.

Statement (iv). For view point $p = (a, b, 0, 0) \in \mathbb{R}^4$ where $a, b \neq 0$, it is easily seen that $j^3\varphi_{p,f} \sim S$ or B in the similar way to the above. Put $p = (a, 0, 0, 0) \in \mathbb{R}^4$ where $a \neq 0$ and we get

$$j^3\varphi_{p,f} \sim_{\mathcal{A}^3} (x, xy - \frac{a_{30}}{a}y^3, b_{21}xy^2 - \frac{b_{30}}{a}y^3).$$

Statement (iv) for $p \in \mathbb{R}^4$ naturally follows, and the case p at infinity is similar. □

Based on Proposition 2.1, we stratify the 3-jet space of Monge forms into strata with codimension at most 2 as in Table 4. In the next section we give simple normal forms of 4-jets of Monge forms which represent each stratum in Table 4 by projective transformations.

Name	Type of 2-jet	Condition	cod	Proj.
Π_E	(x^2-y^2, xy)	$-$	0	$-$
Π_S	(x^2, y^2)	$a_{03}\cdot b_{30}\neq 0$	0	S
Π_B		$b_{30}=0,\ a_{03}\neq 0$	1	S, B
Π_{2B}		$a_{03}=b_{30}=0$	2	B
Π_H	(x^2, xy)	$a_{03}\neq 0$	1	H
Π_P		$a_{03}=0,\ a_{12}\cdot b_{03}\neq 0$	2	P
Π_I^+	$(x^2+y^2, 0)$	$-$	2	S, B
Π_I^-	$(xy, 0)$	$b_{30}\neq 0$	2	S, B, H

TABLE 4. Strata of codimension ≤ 2 in the space of 3-jets of Monge forms corresponding to $\mathcal{A}^3$-types of central projections from view points on asymptotic lines. Surface germs of Π_E-type do not have asymptotic lines.

Remark 2.1. By taking higher order terms of the simple normal forms in the Table 1, we can consider a finer stratification of the space of Monge forms which corresponds to $\mathcal{A}$-types of central projections as Mond did in [15]. For instance, we take the Monge form of the Π_S-type in Table 1 and write $f = (x^2+y^3+\sum_{i+j\geq 4} a_{ij}x^iy^j, y^2+\alpha x^3+\sum_{i+j\geq 4} b_{ij}x^iy^j)$ with $\alpha, a_{ij}, b_{ij}\in\mathbb{R}$, $\alpha\neq 0$ and $a_{40}=b_{04}=0$. Then the condition $a_{31}, b_{13}\neq 0$ determines the proper stratum of the Π_S-stratum where the $\mathcal{A}$-types of the central projection can be determined as the regular, crosscap, S_1 or S_2-type depending on the position of the view points (See [15]).

3. The classification of Monge forms by projective transformations and proof of Theorem 1

In this section we consider a classification of jets of Monge-forms of generic surfaces by projective transformations based on the stratification in Table 4. The projective linear group $PGL(5)$ is defined as the quotient space $GL(5)/\sim$, where $A\sim A'$ if $\exists\lambda\in\mathbb{R}$ such that $A=\lambda A'$. To consider the action on $V_\ell\times V_\ell$ (ℓ-jet-spaces of Monge-forms), we define the following subgroup

$$G(5) := \{\Psi\in PGL(5)\mid \Psi(0)=0,\ \ \Psi(W)=W\}$$

of $PGL(5)$, where $0=[0;0;0;0;1]$ is the origin and W is the xy-plane in $\mathbb{R}^4$. Thus $G(5)$ form a 16-dimensional subgroup of $PGL(5)$ and acts on $V_\ell\times V_\ell$.

Let $f=(f_1,f_2)$ and $g=(g_1,g_2)$ be Monge forms of surface-germs at the origin. We say that the k-jets of these Monge forms are *projectively equivalent* and write $j^kf\sim j^kg$ if there exists $\Psi\in G(5)$ which transforms one to the other. Remark that $\mathcal{A}^\ell$-types of central projections of jets of smooth surfaces are invariant under projective transformations of surfaces, that is, $j^\ell\varphi_{p,f}\sim_{\mathcal{A}^\ell} j^\ell\varphi_{\Phi(p),\Phi(f)}$ from view point p and $\Phi\in G(5)$.

In this paper we check the equivalence of jets of Monge forms in the following way. With the coordinate (x, y, z, w) of $\mathbb{R}^4$, a projective transformation $\Psi \in G(5)$ is regarded locally as a diffeomorphism germ $\mathbb{R}^4, 0 \to \mathbb{R}^4, 0$ given by

$$\Psi(x,y,z,w) = \left(\frac{q_1(x,y,z,w)}{p(x,y,z,w)}, \frac{q_2(x,y,z,w)}{p(x,y,z,w)}, \frac{q_3(x,y,z,w)}{p(x,y,z,w)}, \frac{q_4(x,y,z,w)}{p(x,y,z,w)} \right),$$

where $q_i = q_{i1}x + q_{i2}y + q_{i3}z + q_{i4}w$, for $i = 1, 2$, $q_j = q_{j3}z + q_{j4}w$, for $j = 3, 4$ and $p = 1 + p_1x + p_2y + p_3z + p_4w$. Define

$$F_1(x,y,z,w) = \tfrac{q_3}{p} - f_1(\tfrac{q_1}{p}, \tfrac{q_2}{p})$$
$$F_2(x,y,z,w) = \tfrac{q_4}{p} - f_2(\tfrac{q_1}{p}, \tfrac{q_2}{p}).$$

Then

$$F_1(x, y, g_1, g_2) = F_2(x, y, g_1, g_2) = o(k)$$

where o is Landau's symbol implies $j^k f \sim j^k g$.

Hence, to check the equivalence, we have to solve algebraic equations $F_1 = F_2 = o(k)$ in terms of q_is and p_is for a given Monge form $f = (f_1, f_2)$ and some simplified normal form $g = (g_1, g_2)$. Recall that we already have a stratification of the 3-jet space of Monge forms induced from the $\mathcal{A}^3$-stratification as in Table 4, hence our task is to find a simple normal form of each stratum by projective equivalence. We begin with simplifying 2-jets of Monge forms and then deal with higher jets. However we stop this process with the 4-jets, since the dimension of $G(5)$ which act on the jet space of Monge forms is just 16 and it does not give so good normal forms for higher jets.

3.1. **2-jet.** We first deal with the classification of 2-jets of Monge-forms. In the 2-jet space, the condition $F_1 = F_2 = o(2)$ for any $j^2 f, j^2 g \in V_2 \times V_2$ gives equations of just $q_{i1}, q_{i2}, q_{j3}, q_{j4}$ with $i = 1, 2$ and $j = 3, 4$, and the classification by projective transformations is reduced to the classification of $V_2 \times V_2 \subset J^2(2, 2)$ by the natural action of $\mathcal{G} = GL(2, \mathbb{R}) \times GL(2, \mathbb{R})$. The $\mathcal{G}$-orbits are classified in [9] described as Table 2. We classify now the higher jets of germs with a 2-jet as in Table 2.

3.2. **Elliptic case.** Suppose that $j^2 f = (x^2 - y^2, xy)$ and write

$$j^3(f_1, f_2) = (x^2 - y^2 + \sum_{i+j=3} a_{ij} x^i y^j, xy + \sum_{i+j=3} b_{ij} x^i y^j)$$

where $a_{ij}, b_{ij} \in \mathbb{R}$. The following equivalence

$$j^3(f_1, f_2) \sim (x^2 - y^2 + y^2\phi_1, xy),$$

is given by projective transformation Φ with

$$q_1 = x + b_{03}z + (-a_{31} + b_{12} - b_{30})w, \;\; q_2 = y - b_{30}z + (-b_{21} + b_{03} + a_{30})w,$$
$$q_3 = z, \;\; q_4 = w, \;\; p = 1 + (a_{30} + 2b_{03})x + (2b_{12} - a_{21})y.$$

Here ϕ_k means homogeneous polynomials of degree k. Consider

$$j^4(f_1, f_2) = (x^2 - y^2 + y^2\phi_1 + \sum_{i+j=4} c_{ij} x^i y^j, xy + \sum_{i+j=4} d_{ij} x^i y^j)$$

where $c_{ij}, d_{ij} \in \mathbb{R}$, then

$$j^4(f_1, f_2) \sim (x^2 - y^2 + y^2(\phi_1 + \phi_2), xy + \phi_4),$$

by Φ with $q_1 = x, \;\; q_2 = y, \;\; q_3 = z, \;\; q_4 = w, \;\; p = 1 + c_{40}z + c_{31}w$.

3.3. **Hyperbolic case.** Suppose that $j^2f = (x^2, y^2)$ and write

$$j^3(f_1, f_2) = (x^2 + \sum_{i+j=3} a_{ij}x^iy^j, y^2 + \sum_{i+j=3} b_{ij}x^iy^j)$$

where $a_{ij}, b_{ij} \in \mathbb{R}$. The following equivalence

$$j^3(f_1, f_2) \sim (x^2 + a_{03}y^3, y^2 + b_{30}x^3)$$

is given by projective transformation Φ with

$$q_1 = x + \tfrac{1}{2}(-a_{30} + b_{12})z - \tfrac{1}{2}a_{12}w, \;\; q_2 = y - \tfrac{1}{2}b_{21}z + \tfrac{1}{2}(a_{21} - b_{03})w,$$
$$q_3 = z, \;\; q_4 = w, \;\; p = 1 + b_{12}x + a_{21}y.$$

We can eliminate more two coefficients in 4-jet. Put

$$j^4(f_1, f_2) = (x^2 + a_{03}y^3 + \sum_{i+j=4} c_{ij}x^iy^j, y^2 + b_{30}x^3 + \sum_{i+j=4} d_{ij}x^iy^j)$$

where $c_{ij}, d_{ij} \in \mathbb{R}$, then

$$j^4(f_1, f_2) \sim (x^2 + a_{03}y^3 + y\phi_3, y^2 + b_{30}x^3 + x\psi_3),$$

by Φ with $q_1 = x, \;\; q_2 = y, \;\; q_3 = z, \;\; q_4 = w, \;\; p = 1 - c_{40}z - d_{04}w$. Here ϕ_3 and ψ_3 means homogeneous polynomials of degree 3.

Then

$$j^4(f_1, f_2) \sim \begin{cases} (x^2 + y^3 + y\phi_3, y^2 + \alpha x^3 + x\psi_3), \; \alpha \in \mathbb{R}^* & \text{if } a_{03}, b_{30} \neq 0; \\ (x^2 + y^3 + y\phi_3, y^2 + x\psi_3) & \text{if } a_{03} \neq 0 \text{ and } b_{30} = 0; \\ (x^2 + y\phi_3, y^2 + x\psi_3) & \text{if } a_{03} = b_{30} = 0. \end{cases}$$

3.4. **Parabolic case.** Suppose that $j^2f = (x^2, xy)$ and write

$$j^3(f_1, f_2) = (x^2 + \sum_{i+j=3} a_{ij}x^iy^j, xy + \sum_{i+j=3} b_{ij}x^iy^j)$$

where $a_{ij}, b_{ij} \in \mathbb{R}$. It is easy to show that

$$j^3(f_1, f_2) \sim (x^2 + a_{12}xy^2 + a_{03}y^3, xy + \bar{b}_{12}xy^2 + b_{03}y^3)$$

where $\bar{b}_{12} = b_{12} - \frac{1}{2}a_{21}$. If $a_{03} \neq 0$, then

$$j^3(f_1, f_2) \sim (x^2 + (a_{12} + 3b_{03})xy^2 + a_{03}y^3, xy)$$

with the equivalence given by Ψ with

$$q_1 = x - \tfrac{(-\bar{b}_{12}a_{03} + 3a_{12}b_{03} + 3b_{03}^2)}{a_{03}}w,$$
$$q_2 = \tfrac{b_{03}}{a_{03}}x + y + \tfrac{b_{03}^2(a_{12}b_{03} - a_{03}\bar{b}_{12})}{a_{03}^3}z - \tfrac{b_{03}(2b_{03}^2 + \bar{b}_{12}a_{03})}{a_{03}^2}w, \;\; q_3 = z,$$
$$q_4 = \tfrac{b_{03}}{a_{03}}z + w, \;\; p = 1 + \tfrac{b_{03}^2(a_{12} + b_{03})}{a_{03}^2}x - \tfrac{(-2\bar{b}_{12}a_{03} + 4a_{12}b_{03} + 3b_{03}^2)}{a_{03}}y.$$

Then the 4-jet can be written in the form

$$j^4(f_1, f_2) \sim (x^2 + \beta xy^2 + y^3 + y\phi_3, xy + x\psi_3),$$

where $\beta = \frac{(a_{12} + 3b_{03})}{a_{03}^{3/2}}$, ϕ_3 and ψ_3 mean homogeneous polynomials of degree 3.

If $a_{03} = 0$ but $a_{12} \neq 0$ we obtain

$$j^3(f_1, f_2) \sim (x^2 + xy^2, xy + \gamma y^3)$$

with the projective transformation Ψ given by

$$q_1 = a_{12}x + a_{12}b_{12}w, \;\; q_2 = y, \;\; q_3 = a_{12}^2 z,$$
$$q_4 = a_{12}w, \;\; p = 1 + 2b_{12}y,$$

where $\gamma = \frac{b_{03}}{a_{12}}$. If we put

$$j^4(f_1, f_2) = (x^2 + xy^2 + \sum_{i+j=4} a_{ij}x^i y^j, xy + \beta y^3 + \sum_{i+j=4} b_{ij}x^i y^j),$$

then $\gamma \neq 0$ leads to

$$j^4(f_1, f_2) \sim (x^2 + xy^2 + \lambda y^4, xy + \gamma y^3 + \phi_4)$$

by a projective transformation Ψ with

$$\begin{gathered} q_1 = x + \tfrac{1}{2}(-q_{21}^2 + p_1)z + (3\gamma q_{21} - 3q_{21})w, \\ q_2 = y + q_{21}x + \tfrac{1}{2}(-2\gamma q_{21}^3 + q_{21}^3 + p_1 q_{21})z + \tfrac{1}{2}(-q_{21}^2 + p_1)w, \\ q_3 = z, \;\; q_4 = q_{21}z + w, \;\; p = 1 + p_1 x - (-6\gamma q_{21} - 4q_{21})y + p_3 z + p_4 w, \end{gathered}$$

where $p_1 = \frac{1}{\Lambda^2}\xi_1$, $p_3 = \frac{1}{\Lambda^4}\xi_2$, $p_4 = \frac{1}{\Lambda^3}\xi_3$, $q_{21} = -\frac{a_{13}}{\Lambda}$, ξ_i are combinations of the coefficients of 4-jet and $\Lambda = 6\gamma^2 + 4\lambda - 15\gamma + 5 \neq 0$. If $\Lambda = 0$ the terms of $j^4(f^1, f^2)$ of order 4 can not be removed. The ϕ_4 is a homogeneous polynomials of degree 4.

3.5. **Inflection case.** Suppose that $j^2 f = (x^2 + y^2, 0)$ and write

$$j^3(f_1, f_2) = (x^2 + y^2 + \sum_{i+j=3} a_{ij}x^i y^j, \sum_{i+j=3} b_{ij}x^i y^j).$$

Let $b_{30} - b_{12} \neq 0$. It follows that

$$j^3(f_1, f_2) \sim (x^2 + y^2 + k_1 x^2 y, \phi_3)$$

by Ψ with

$$\begin{gathered} q_1 = x, \;\; q_2 = y, \;\; q_3 = z + (a_{30} - a_{12})w, \;\; q_4 = (b_{30} - b_{12})w, \\ p = 1 - \tfrac{(a_{30}b_{12} - a_{12}b_{30})}{(b_{30} - b_{12})}x - \tfrac{(a_{30}b_{03} - a_{12}b_{03} - a_{03}b_{30} + a_{03}b_{12})}{(b_{30} - b_{12})}y. \end{gathered}$$

Here, k_1 is scalar constant. Now, we take

$$j^4(f_1, f_2) = (x^2 + y^2 + k_1 x^2 y + \sum_{i+j=4} c_{ij}x^i y^j, \phi_3 + \sum_{i+j=4} d_{ij}x^i y^j),$$

where $c_{ij}, d_{ij} \in \mathbb{R}$, then it follows that

$$j^4(f_1, f_2) \sim (x^2 + y^2 + k_1 x^2 y + y\psi_3, \phi_3 + \phi_4)$$

by Ψ with $q_1 = x, \;\; q_2 = y, \;\; q_3 = z, \;\; q_4 = w, p = 1 + c_{40}z$. Here ϕ_k and ψ_k means homogeneous polynomials of degree k.

Next, suppose

$$j^3(f_1, f_2) = (xy + \sum_{i+j=3} a_{ij}x^i y^j, \sum_{i+j=3} b_{ij}x^i y^j).$$

If $b_{03} \neq 0$, then

$$j^3(f_1, f_2) \sim (xy + k_2 x^3, \phi_3)$$

by Ψ with

$$\begin{gathered} q_1 = x, \;\; q_2 = y, \;\; q_3 = z + (a_{21} + a_{03})w, \;\; q_4 = b_{03}w, \\ p = 1 - \tfrac{(a_{21}b_{03} - a_{03}b_{21})}{b_{03}}x - \tfrac{(a_{21}b_{03} - a_{03}b_{12})}{b_{03}}y. \end{gathered}$$

The k_2 is a scalar constants. Finally, we consider

$$j^4(f_1, f_2) = (xy + k_2 x^3 + \sum_{i+j=4} c_{ij}x^i y^j, \phi_3 + \sum_{i+j=4} d_{ij}x^i y^j),$$

where $c_{ij}, d_{ij} \in \mathbb{R}$. Thus, it follows that

$$j^4(f_1, f_2) \sim (xy + k_2x^3 + \bar{\xi}_4, \phi_3 + \phi_4)$$

by Ψ with $q_1 = x,\ q_2 = y,\ q_3 = z,\ q_4 = w,\ p = 1+c_{22}z$. The ϕ_k means homogeneous polynomials of degree k and $\bar{\xi}_4$ is a homogeneous polynomials of degree 4 without the term x^2y^2.

References

[1] V. I. Arnold, *Catastrophe Theory*, 3rd edition, Springer (2004).

[2] V. I. Arnold, V. V. Goryunov, O. V. Lyashko, V. A. Vasil'ev, *Singularity Theory II, Classification and Applications*, Encyclopaedia of Mathematical Sciences Vol. 39, Dynamical System VIII (V. I. Arnold (ed.)), (translation from Russian version), Springer-Verlag (1993).

[3] J. W. Bruce, Projections and reflections of generic surfaces in $\mathbb{R}^3$. *Math. Scand.* 54 (1984), 262–278.

[4] J. W. Bruce, P. J. Giblin and F. Tari. Families of surfaces: height function, Gauss maps and duals. In *Real and Complex singularities (São Carlos, 1994)*, 148–178. Pitman Res. Notes Math. Ser. 333, Longman, Harlow, 1995.

[5] J. W. Bruce and A. C. Nogueira, Surfaces in $\mathbb{R}^4$ and duality. *Quart. J. Math. Oxford Ser.* Ser. (2), 49 (1998), 433–443.

[6] J. W. Bruce and F. Tari, Families of surfaces in $\mathbb{R}^4$. *Proc. Edinb. Math. Soc.* 45 (2002), 181–203.

[7] J. L. Deolindo-Silva. Cr-invariants for surfaces in $\mathbb{R}^4$. Preprint, 2015.

[8] J. L. Deolindo-Silva, Y. Kabata, T. Ohmoto, H. Sano. Projective classification of jets of surfaces in 3-space, preprint, (2015).

[9] C. G. Gibson, Singular points of smooth mappings, Pitman Research Notes in Mathematics, vol. 25 (1979).

[10] V.V. Goryunov, Singularities of projections of complete intersections, *J. Soviet Math.* 27 (1984), 2785–2811.

[11] Y. Kabata, Recognition of plane-to-plane map-germs, preprint (2015), ArXiv: 1503.08544.

[12] E. E. Landis, Tangential singularities, Funt. Anal. Appl. 15 (1981), 103–114 (translation).

[13] J. A. Little, On the singularities of submanifolds of heigher dimensional Euclidean space. *Annli Mat. Pura et Appl.* (4A) 83 (1969), 261–336.

[14] D. K. H. Mochida, M. C. Romero-Fuster and M. A. S. Ruas, The geometry of surfaces in 4-space from a contact viewpoint. *Geometria Dedicata* 54 (1995), 323–332.

[15] D. M. Q. Mond, Classification of certain singularities and applications to differential geometry. Ph.D. thesis, The University of Liverpool, 1982.

[16] D. M. Q. Mond, On the classification of germs of maps from $\mathbb{R}^2$ to $\mathbb{R}^3$. *Proc. London Math. Soc.* 50 (1985), 333–369.

[17] D. M. Q. Mond. Some remarks on the geometry and classification of germs of maps from surfaces to 3-space. *Topology.* 26 (3) (1987), 361–383.

[18] J. J. Nuño-Ballesteros and F. Tari, Surfaces in $\mathbb{R}^4$ and their projections to 3-spaces. *Roy. Proc. Edinburgh Math. Soc.* 137A (2007), 1313–1328.

[19] R. Oset-Sinha, F. Tari, Projections of surfaces in $\mathbb{R}^4$ to $\mathbb{R}^3$ and the geometry of their singular images. *Rev. Mat. Iberoam.* 31, no. 1, 33-50, 2015.

[20] O. A. Platonova, Singularities of the mutual disposition of a surface and a line, Uspekhi Mat. Nauk, 36:1 (1981), 221–222.

[21] O. A. Platonova, Projections of smooth surfaces, *J. Soviet Math.* 35 no.6 (1986), 2796–2808